

Enriched eggs as a source of n-3 polyunsaturated fatty acids for humans

Gordana Kralik¹, Zlata Kralik¹, Eva Strakova², Manuela Grčević¹, Danica Hanžek¹

¹Josip Juraj Strossmayer University of Osijek, Faculty of Agriculture in Osijek, Department of Special Zootechniques, Osijek, Croatia (gkralik@pfos.hr)

²University of Veterinary and Pharmaceutical Science Brno, Faculty of Veterinary Hygiene and Ecology, Department of Nutrition, Animal Husbandry and Animal Hygiene, Brno, Czech Republic

The aim of the research was to enrich egg with n-3 polyunsaturated fatty acids by using plant oils and fish oil as dietary supplements in laying hens' feed. The focus was put on the effect of the daily consumption of 100 g of egg yolks, i.e. 100 g of egg mass, on the human health. The 1st group of laying hens was fed a diet containing soybean and fish oil, and the 2nd group was given feed containing a combination of linseed, rapeseed, soybean, and fish oils. Eggs laid by the 2nd group contained 4.73% α -linolenic acid, 0.20% eicosapentaenoic acid and 2.37% docosahexaenoic acid (% of total fatty acids in yolk lipids, $P < 0.001$), which marks an increase of $\times 4.04$ for α -linolenic acid, $\times 3.33$ for eicosapentaenoic acid, and $\times 1.75$ for docosahexaenoic acid compared to eggs laid by the 1st group. Total n-3 polyunsaturated fatty acids in eggs of the 2nd group were $\times 2.8$ higher than in the 1st group. Calculated per 100 g of eggs of the 2nd group, the intake for the human body corresponds to 435 mg α -linolenic acid, 18.43 mg eicosapentaenoic acid, and 218.2 mg docosahexaenoic acid.

Key words:

deposition, ALA, EPA, DHA, consumption

1. Introduction

Nutritionists recommend consumption of eggs enriched with n-3 polyunsaturated fatty acids (n-3 PUFA), having in mind the necessity of preventing various diseases and strengthening the immune system in humans. Benefit for human health is estimated on the basis of n-3 PUFA content in egg yolk lipids, as well as on the ratio of n-6 / n-3 PUFA (Okuyama et al. 1997; Simopoulos 2000). Omega-3 fatty acids prevent the occurrence of rheumatoid arthritis and cancer (Aydin and Dogan 2010), as well as cardiovascular diseases (Van Elswik 1997; Raes et al. 2002; Pita et al. 2010), and have positive effects on the immune system (He et al. 2007). Rašić et al. (2014) determined that consumption of eggs enriched with n-3 PUFA over a period of three weeks had positive effect on microvascular reactivity, on the lowering of blood pressure and triglycerides, so they concluded that omega-3 eggs could be efficient in preventing cardiovascular diseases. Laying hens have limited ability to convert alpha-linolenic acid (ALA) into eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) in their organism, and it is recommended that these fatty acids are taken in through feed, because in that way the enrichment of eggs with n-3 PUFA was more effective (Kralik 2007; Škrtić 2007, 2008). Oils (linseed, fish, rapeseed, soybean) are generally used as a source of energy from the laying hens' feed. Various studies have shown that the type of oil could have a significant impact on the profile of fatty

The article was originally published in the journal „Acta Veterinaria Brno“ 2017, 86: 293-301; <https://doi.org/10.2754/avb201786030293>. The reprint permission granted by the authors.

acids in egg yolks (Rowghani et al. 2007; Aydin and Dogan 2010; Herkel et al., 2016). Fish oil is a rich source of essential n-3 PUFA, but if it is used in large amounts in feeding mixtures for poultry, then it has negative effect on organoleptic characteristics of meat and eggs (Scaife et al. 1994; Scheideler 1997). Rapeseed and linseed oils are well known sources of ALA that convert to long chain n-3 polyunsaturated acids, such as EPA and DHA, through the elongation and desaturation processes (Yang et al. 2000).

The aim of this research was to enrich eggs with n-3 PUFA, especially with EPA and DHA, by using a combination of linseed, rapeseed, soybean and fish oils in laying hens' diet in the amount of 5%. Combination of different oils is interesting because of the interaction of individual fatty acids in the metabolic processes, as well as because of the impact on altering the n-6/n-3 PUFA ratio in yolk lipids. The control group of laying hens was given a diet containing 4.5% soybean oil and 0.5% fish oil. This paper aimed to investigate how and in what concentration the sources of ALA, EPA and DHA from oils, i.e. from feed, could influence their deposition in yolk lipids of table eggs. Furthermore, the intention was to determine effects of the stated n-3 PUFA by consumption of 100 g of egg yolks, i.e. 100 g of egg mass of enriched eggs and to compare them with conventional eggs.

2. Materials and Methods

The research was carried out on the Lohmann Classic laying hens aged 30-34 weeks. Laying hens were divided into two groups (control and experimental). Each group consisted of five cages, and in each cage there were 40 hens (on the area of 750 cm²/head), i.e. there were 200 hens in the 1st group and 200 hens in the 2nd group. Laying hens were fed diets of isoprotein and isocaloric composition during a period of four weeks (Table 1).

Table 1. Ingredients and content of diets

Ingredients, %	1 st group	2 nd group
Corn	49.80	48.83
Soybean cake	26.13	25.62
Sunflower cake	3.91	5.00
Dehydrated alfalfa	2.00	2.00
Livestock yeast	0.50	0.50
Limestone	10.18	10.57
Monocalcium phosphate	1.51	1.51
Salt	0.32	0.32
Methionine	0.15	0.15
Oil	5.00 ¹	5.00 ²
Premix ³	0.50	0.50
Total	100.00	100.00
Calculative values, %		
Crude protein	18,0	18,0
Crude fat	7.4	7.4
Crude fibers	4.0	3.8
Ash	13.7	13.3
ME MJ/kg	11.5	11.5

¹ soybean oil 4.5%, fish oil 0.5%; ² oil combination: linseed oil 1%, fish oil 0.75%, rapeseed oil 2%, soybean oil 1.25%; ³Premix contains in 1 kg: vitamin A 200,000 UI, vitamin D₃ 500,000 UI, vitamin E 10,000 mg, vitamin K₃ 600 mg, vitamin B₁ 400 mg, vitamin B₂ 1,000 mg, vitamin B₆ 1,000 mg, vitamin B₁₂ 3,000 µg, vitamin C 4,000 mg, vitamin H 12 mg, vitamin B₃ 8,000 mg, vitamin B₅ 2,400 mg, vitamin B₉ 150 mg, vitamin B₄ 100,000 mg, iodine 200 mg, manganese 18,000 mg, zinc 14,000 mg, cobalt 30 mg, iron 12,000 mg, copper 1,600 mg, selenium inorganic 50 mg, calcium 238 g, phytase 100,000 FYT, canthaxanthin 500 mg, beta-apo-beta carotene acid 300 mg, antioxidant (butylated hydroxytoluene) 20,000 mg

Apart of other feed ingredients, the 1st group of hens was given diets containing 4.5% of soybean oil and 0.5% of fish oil (total 5%), whereas the 2nd group was fed diets composed of a combination of the following oils: linseed 1%, fish 0.75%, 2% rapeseed 2%, and soybean 1.25%, which amounts to the same total of 5%. Laying hens were administered feed and water *ad libitum*. During the experiment, the lighting regime was applied according to technological standards. Eggs used for the analysis were collected on the last day of the experiment. Two eggs were collected from each cage (total n = 10 pcs).

Profile of fatty acids in egg yolk lipids was determined according to the method of Csapo et al. (1986) by using Chrompack CP-9000 gas chromatograph equipped with flame ionization detector. The operating conditions of the gas chromatography were as follows: 0.35 grams of dried egg-yolk was weighed into a flask, 8 cm³ concentrated hydrochloric acid was added and it was boiled for 60 minutes. After cooling down 7 cm³ ethanol was added, then 15 cm³ diethylether following one minute of shaking. The next extraction was with 15 cm³ benzene (b.p. <60 °C). After phase separation, the organic phase that contains about 150-200 mg fat was separated and evaporated under vacuum on a rotadest. Then 4 cm³ 0.5 M sodium-hydroxide in methanol was added, and boiled in a water bath for 5 minutes. Next 4 cm³ 14% boron-trifluoride in methanol was added and boiled for 3 minutes following the addition of 4 cm³ n-hexane. It was boiled for one minute after which the level of the organic phase was brought to the neck of the flask with saturated sodium-chloride solution. When the phases were separated, samples were taken for analysis from the organic phase, and it was dried on sodium sulphate. The fatty acid methyl esters (FAMES) were separated on a 100 m x 0.25 mm wall coated open tubular (WCOT) column equipped with a CP-SIL 88 (FAME) stationary phase. The quantitation of FAMES was obtained with a flame ionisation detector (FID) at 270 °C. The temperature of the splitter injector was 270 °C, the carrier gas was helium with the head pressure of 235 kPa. The oven was temperature programmed from 140 °C (10 min) with 10 °C/min increase up to 235 °C (26 min). The injected volume varied between 0.5 and 2 µl. The instrument was a Chrompack CP 9000 gas chromatograph. Portions of saturated fatty acids (SFA) and monounsaturated fatty acids (MUFA), ALA, as well as EPA and DHA acids show as a percentage of total fatty acids contained in yolk lipids. The influence of different feeding treatments (oils) was determined by t test. Research results were processed by Statistica v. 8.0.

3. Results

Contents of fatty acids in linseed, fish, rapeseed and soybean oils (% of total FA) are presented in the Table 2. Linseed oil contains 54.82% ALA. Rapeseed oil and soybean oil contain the most monounsaturated fatty acids (51.23% and 64.82%, respectively), of which oleic acid (OA, 42.86% and 63.89%, respectively) is the most represented. Rapeseed and soybean oils also contain linoleic acid (LA, 19.75% and 20.55%, respectively), and ALA (14.48% and 6.63%, respectively). Fish oil is rich in saturated fatty acids (SFA, 38.90%), and EPA (7.15%) and DHA (18.34%) are represented from the n-3 PUFA group.

Table 2. Profile of fatty acids in oils (% of total fatty acids)

Fatty acid	Linseed oil	Fish Oil	Rapeseed oil	Soybean oil
Myristic acid (C14:0)	0.11	6.14	1.41	0.06
Pentadecanoic acid (C15:0)	0.03	1.98	0.52	0.02
Palmitic acid (C16:0)	6.81	21.92	7.66	4.70
Heptadecanoic acid (C17:0)	0.16	1.31	0.32	0.04
Stearic acid (C18:0)	3.83	5.90	2.82	2.34
Arachidic acid (C20:0)	0.17	1.03	0.46	0.49
Behenic acid (C22:0)	0.13	0.62	0.00	0.24
ΣSFA*	11.23	38.90	13.20	7.89
Palmitoleic acid (C16:1)	0.05	5.10	1.40	0.20
Heptadecenoic acid (C17:1)	0.00	1.27	0.00	0.00
Oleic acid (C18:1n9c)	18.78	17.34	42.86	63.89
Eicosenoic acid (C20:1n9)	0.22	2.21	4.38	0.86
Erucic acid (C22:1n9)	0.00	0.15	2.59	0.04
ΣMUFA**	19.04	26.07	51.23	64.82
Linoleic acid (C18:2n6)	14.57	3.65	19.75	20.55
γ-linolenic acid (C18:3n6)	0.12	0.48	0.49	0.00
Eicosadienoic acid (C20:2)	0.06	0.68	0.43	0.03
Eicosatrienoic (C20:3n6)	0.00	0.15	0.00	0.00
Arachidonic acid (C20:4n6)	0.00	1.41	0.29	0.00
Docosadienoic acid (C22:2n6)	0.00	0.63	0.00	0.08
Σn-6 PUFA***	14.75	7.00	20.96	20.66
α-linolenic acid (C18:3n3)	54.82	2.45	14.48	6.63
Eicosatrienoic acid (C20:3n3)	0.15	0.10	0.13	0.00
Eicosapentaenoic acid (C20:5n3)	0.00	7.15	0.00	0.00
Docosahexaenoic acid (C22:6n3)	0.00	18.34	0.00	0.00
Σn-3 PUFA	54.97	28.04	14.61	6.63
Σn-6 PUFA / Σn-3 PUFA	0.27	0.25	1.43	3.12

*SFA - saturated fatty acids; **MUFA - monounsaturated fatty acids; ***PUFA - polyunsaturated fatty acids

The ratio n-6/n-3 PUFA in oils is very narrow, ranging from 0.27 (for linseed oil) to 3.12 (for soybean oil). Fatty acid profile in laying hens' feed depends on the type and concentration of supplemented oils (Table 3).

The greatest difference between laying hens diets was noticed in the content of n-6 PUFA and n-3 PUFA. The 1st group being fed diet supplemented with 4.5% soybean oil and 0.5% fish oil contained 42.57% of n-6 PUFA and only 3.34% of n-3 PUFA, so the ratio of n-6/n-3 PUFA was 12.75. Diet given to the 2nd group contained 37.28% of n-6 PUFA and 17.40% of n-3 PUFA and their ratio was 2.14. The most represented fatty acid in n-6 PUFA group in both laying hens' diets was LA (41.49% and 35.98%, respectively). In total n-3 PUFA, ALA was present in significant amounts in both diets (3.12% and 13.36%, respectively). It should be pointed out that the content of DHA in the diet of the 1st group was 68.5 times higher than in the diet given to the 2nd group.

Table 3. Profile of fatty acids in laying hens' diet (% of total fatty acids)

Fatty acid	1 st group	2 nd group
Myristic acid (C14:0)	2.27	1.69
Pentadecanoic acid (C15:0)	0.09	0.25
Palmitic acid (C16:0)	13.07	12.73
Heptadecanoic acid (C17:0)	0.12	0.13
Stearic acid (C18:0)	3.29	3.19
Arachidic acid (C20:0)	0.55	0.41
Behenic acid (C22:0)	0.79	0.70
Lignoceric acid (C24:0)	0.26	0.17
ΣSFA*	20.43	19.27
Palmitoleic acid (C16:1)	0.20	0.51
Oleic acid (C18:1n9c)	33.02	24.97
Eicosenoic acid (C20:1n9)	0.40	0.51
Erucic acid (C22:1n9)	0.04	0.06
ΣMUFA**	33.66	26.05
Linoleic acid (C18:2n6)	41.49	35.98
γ-linolenic acid (C18:3n6)	0.29	0.32
Eicosadienoic acid (C20:2)	0.29	0.30
Eicosatrienoic (C20:3n6)	0.10	0.13
Arachidonic acid (C20:4n6)	0.14	0.07
Docosadienoic acid (C22:2n6)	0.26	0.48
Σn-6 PUFA***	42.57	37.28
α-linolenic acid (C18:3n3)	3.12	13.36
Eicosatrienoic acid (C20:3n3)	0.09	0.15
Eicosapentaenoic acid (C20:5n3)	0.09	1.16
Docosahexaenoic acid (C22:6n3)	0.04	2.74
Σn-3 PUFA	3.34	17.40
Σn-6 PUFA/Σn-3 PUFA	12.75	2.14

*SFA – saturated fatty acids; **MUFA - monounsaturated fatty acids; ***PUFA - polyunsaturated fatty acids

The Table 4 overviews the profile of fatty acids (% of total FA) in yolk lipids of the 1st and 2nd group. In yolk lipids, eggs laid by the 2nd group contained statistically significantly less ΣSFA than the 1st group (31.33% : 34.72%, $P < 0.001$). Similar results were obtained by Kralik et al. (2008) in their previous research of usage of fish oil in production of table eggs. Eggs produced in the 2nd group contained significantly ($P < 0.05$) less palmitoleic acid, and more eicosenoic acid. Differences in total content of MUFA and n-6 PUFA between eggs of both groups were not significant ($P > 0.05$).

Table 4. Fatty acids profile in lipids of egg yolk (% of total fatty acids, n=10)

Fatty acid	1 st group $\bar{x} \pm \text{sd}$	2 nd group $\bar{x} \pm \text{sd}$	P-value
Myristic acid (C14:0)	0.34 ^a \pm 0.03	0.29 ^b \pm 0.03	0.005
Pentadecanoic acid (C15:0)	0.20 \pm 0.03	0.19 \pm 0.04	0.688
Palmitic acid (C16:0)	22.16 ^a \pm 1.13	19.86 ^b \pm 1.58	0.001
Heptadecanoic acid (C17:0)	0.34 \pm 0.07	0.34 \pm 0.04	0.759
Stearic acid (C18:0)	8.29 \pm 1.13	7.69 \pm 1.20	0.265
Arachidic acid (C20:0)	0.17 \pm 0.04	0.17 \pm 0.02	0.928
Heneicosanoic acid (C21:0)	0.12 \pm 0.02	0.13 \pm 0.02	0.426
Tricosanoic acid (C23:0)	3.00 \pm 0.36	2.66 \pm 0.61	0.144
Σ SFA*	34.72 ^a \pm 1.13	31.33 ^b \pm 0.64	<0.001
Myristoleic acid (C14:1)	0.17 \pm 0.04	0.17 \pm 0.03	0.705
Palmitoleic acid (C16:1)	2.43 ^a \pm 0.35	2.00 ^b \pm 0.30	0.008
Elaidic acid (C18:1n9t)	0.02 \pm 0.004	0.02 \pm 0.002	0.151
Oleic acid (C18:1n9c)	38.67 \pm 1.93	38.94 \pm 1.28	0.725
Eicosenoic acid (C20:1n9)	0.22 ^b \pm 0.04	0.27 ^a \pm 0.03	0.002
Erucic acid (C22:1n9)	0.31 ^b \pm 0.04	0.37 ^a \pm 0.05	0.013
Σ MUFA**	41.82 \pm 2.04	41.76 \pm 1.08	0.939
Linoleic acid (C18:2n6)	19.13 \pm 1.87	17.83 \pm 0.83	0.060
γ -linolenic acid (C18:3n6)	0.24 \pm 0.03	0.22 \pm 0.03	0.085
Eicosadienoic acid (C20:2n6)	0.26 \pm 0.08	0.32 \pm 0.08	0.158
Eicosatrienoic acid (C20:3n6)	0.36 \pm 0.05	0.35 \pm 0.05	0.641
Arachidonic acid (C20:4n6)	0.57 \pm 0.07	0.61 \pm 0.05	0.128
Docosadienoic acid (C22:2n6)	0.28 \pm 0.04	0.26 \pm 0.02	0.085
Σ n-6 PUFA***	20.85 \pm 1.91	19.59 \pm 0.83	0.071
α -linolenic acid (C18:3n3)	1.17 ^b \pm 0.15	4.73 ^a \pm 0.21	<0.001
Eicosatrienoic (C20:3n3)	0.02 \pm 0.01	0.03 \pm 0.01	0.115
Eicosapentaenoic acid (C20:5n3)	0.06 ^b \pm 0.02	0.20 ^a \pm 0.03	<0.001
Docosahexaenoic acid (C22:6n3)	1.35 ^b \pm 0.22	2.37 ^a \pm 0.18	<0.001
Σ n-3 PUFA	2.60 ^b \pm 0.18	7.32 ^a \pm 0.23	<0.001
Σ n-6/ Σ n-3 PUFA	8.02 ^a \pm 0.64	2.68 ^b \pm 0.12	<0.001

\bar{x} = arithmetic means; sd = standard deviation; ^{a,b}P < 0.05, P < 0.01 or P < 0.001 ; *SFA – saturated fatty acids; **MUFA – monounsaturated fatty acids; ***PUFA –polyunsaturated fatty acids

Table 5 presents concentrations of ALA, EPA and DHA in egg yolks and in total egg mass of the 1st and 2nd group.

Table 5. Concentration of n-3 PUFA in yolks and in eggs

Fatty acid	1 st group		2 nd group	
	mg/100 g of yolk	mg/100 g of egg	mg/100 g of yolk	mg/100 g of egg
ALA*	254.2	108.0	1102.0	435.0
EPA**	14.10	5.99	46.63	18.43
DHA**	291.7	124.0	552.1	218.2

*ALA - α -linolenic acid; **EPA – eicosapentaenoic acid; *** DHA – docosahexaenoic acid

The content of ALA was 254.2 mg/100 g in egg yolks of the 1st group, and 1102 mg/100 g of the 2nd group. The content of EPA was increased from 14.1 mg/100 g to 46.63 mg/100 g. The content of DHA was also increased from 291.7 to 552.1 mg/100 g of yolk. In comparison to the content of FA in whole eggs of the 1st group, eggs of the 2nd group had ALA increased by x 4.02, EPA by x 3.07, and DHA by x 1.75.

4. Discussion

Our research results regarding the content of n-3 PUFA in oils proved that there were some differences. Linseed oil is rich in ALA (54.85%), fish oil is rich in DHA (18.34%) and EPA (7.15%). Rapeseed oil and soybean oil contain great amounts of oleic acid (42.86% and 63.89%, respectively). As of the n-3 PUFA group, ALA is the only one present in both types of oil. These results are in accordance with those published by Rowghani et al. (2007), Kralik et al. (2008), Antongiovanni et al. (2009) and Salamatdoustnobar et al. (2009), although there is very little fluctuation referring to FA, caused by the origin of oil. In general, plant oils contain less than 25% ALA, with the exception of linseed oil, which is rich in ALA (Hiltunen and Holm 2000; Herkel et al. 2016). ALA in linseed oil varies between 34.1 and 64.4% (Hiltunen and Holm 2000). The time of harvest and plant processing affects the yield and quality of linseed oil (Nykter and Kymäläinen 2006). The n-6 PUFA/n-3 PUFA ratio in linseed oil is 4:1. The diet fed to the 2nd group contained more MUFA than diet of the 1st group (33.66% : 26.05%, respectively), but the most common one was oleic acid (33.02% : 24.07%, respectively). The diet for the 1st group was richer in linoleic acid than the diet composed for the 2nd group (41.49% : 35.98%). At the same time, the 2nd group was given a diet containing more ALA, EPA and DHA than the 1st group (13.36%, 1.16% and 2.74% compared to 3.12%, 0.09% and 0.04%, respectively).

We did not establish significant differences in LA between groups of eggs, although eggs from the 2nd group contained less LA, but slightly more arachidonic acid ($P > 0.05$), which is the opposite to our previous research results (Kralik et al. 2008), as well as to research results of Sari et al. (2002) and Husveth et al. (2003). Beynen (2004) determined that only high concentration of LNA in a diet affected the synthesis of detectable content of EPA in yolks, which was generally lower than the content of DHA. The author also stated that LA from feed incorporated into the yolk more efficiently than ALA.

Eggs from hens of the 2nd group contained more n-3 PUFA than the 1st group ($P < 0.001$). When compared to the 1st group, higher portions of ALA (4.73% : 1.17%), EPA (0.20% : 0.06%), DHA (2.37% : 1.35%) and n-3 PUFA (7.32% : 2.60%) were determined in the yolk lipids of the 2nd group. Although diet given to the 1st group of hens contained smaller amounts of EPA and DHA, these fatty acids were detected in yolk lipids, because laying hens can synthesize EPA and DHA at a limited amount if the diet contains ALA in sufficient amount (Kralik et al. 2008). Increased amounts of EPA and DHA in feed resulted in increased deposition of DHA in the yolks lipids of the 2nd group, which is a result of dietary supplementation with different oils. Similar results were achieved by Bavelaar and Beynen (2004) and Huyghebaert et al. (2007). Kralik et al. (2008) reported that dietary supplementation of laying hens' feed with various combinations of fish and rapeseed oils, instead of soybean oil, resulted in the increased content of n-3 PUFA for x 1.4 to 2.10.

Rowghani et al. (2007) stated that rapeseed oil (3% and 5%) increased the percentage of ALA to 3.43% and 6.02%, respectively, and also affected the increase of DHA. Linseed oil also increased Σ n-3 PUFA to 4.72% and 6.80% when compared to the control (1.43%), which was also found by Herkel et al. (2016). Lewis et al. (2000) determined that supplementation of 7% rapeseed oil affected the increase of ALA by x 1.2% to 6.3%, and the increase of DHA x 4. The authors concluded that three eggs enriched with n-3 PUFA correspond to the amount of n-3

PUFA in one fish meal daily. Bearing in mind that people are more prone to consume eggs than fish, they can thus be an alternative source of n-3 PUFA. According to the Canadian standards, it is recommended to consume eggs enriched with n-3 PUFA by up to 50% of energy in a meal. Samman et al. (2009) stated that omega eggs had significantly lower percentage of myristic and palmitic acid, which resulted in a lower percentage of SFA, but significantly higher percentage of ALA, DHA and total omega-3, which is also the case in our research. Gül et al. (2012) pointed out that dietary supplementation of laying hens' feed with rapeseed oil resulted in increased MUFA, especially of oleic acid, in eggs. In this research, oleic acid in egg yolk was at approximately the same level in both groups (>45%).

The ratio of n-6/n-3 PUFA in yolk lipids was 2.68 in the 2nd group, and 8.02 in the 1st group ($P < 0.001$). Sugano (1996) stated that the optimal ratio of n-6/n-3 PUFA in human nutrition was 3:1, however, some studies in developed industrial countries proved this ratio to range from 10:1 and to 15:1 and more (Simopoulos 2000). Our research proved that the feeding treatment in the 2nd group (combination of plant and fish oils) was very efficient in enriching yolk lipids with n-3 PUFA, and consequently in reducing the ratio of n-6/n-3 PUFA, compared to the feeding treatment with soybean oil and fish oil.

In line with Meluzzi et al. (2000), Kralik et al. (2008) and Herkel et al. (2016), enrichment of eggs with the n-3 PUFA depended on the increased content of n-3 PUFA in hens' diet, and it was connected with the lowering of n-6 PUFA in egg yolks. According to Simopoulos (2000), LA and ALA are very important as precursors of n-6 PUFA and n-3 PUFA. Linoleic acid is metabolized into other n-6 fatty acids, including AA. Alpha-linolenic acid is metabolized into the n-3 fatty acids, such as EPA and DHA. When LA is present at larger amounts, it will inhibit the transformation of ALA into EPA and DHA, and with insufficient amount of LA, lower AA concentration will be created, which was not the case in our research. Our results show that narrowing of n-6/n-3 PUFA ratio and the increase of the n-3 PUFA content in feed can affect the FA profile in yolk lipids directed towards the increased deposition of n-3 PUFA, which was also reported by Beynen (2004), Škrčić et al. (2007), as well as by Kralik et al. (2008). Bavelaar and Beynen (2004) proved a linear relationship between ALA and LA in feed and their contents in yolk lipids.

The recommended daily intake of n-3 PUFA is 0.5% of the total energy in a meal (Anon 1990), which is an equivalent to 1.1-1.5 g FA for adults. Ferrier et al. (1995) calculated that feeding laying hens with diets containing 10% linseed oil, which is rich in ALA, result in production of eggs, where only one egg covers 30% of the n-3 PUFA daily requirements. Such egg contains 264 mg ALA, 10 mg EPA and 82 mg DHA. Jiang et al. (1991) also reported that EPA and DHA could be increased in yolk lipids when the feed was rich in ALA that was converted to EPA and DHA in the laying hens' liver, and synthesized fatty acids were deposited in the egg yolks. That conversion may be inhibited by high LA concentration. Beynen (2004) determined that higher amounts of linseed in feed affected the increased deposition of EPA and DHA in yolk lipids, whereas the feeding of hens with soybean was less efficient. The author concluded that the diet enriched with linseed or soybean influenced the deposition of 141 mg and 98 mg of DHA, unlike in the control group, where egg yolks contained 78 mg of DHA. Favorable effects on human health can be achieved by daily consumption of 0.5 g of n-3 PUFA (Mantzioris 2000). The results of this research proved that 100 g of egg mass from the 2nd group contained 435 mg ALA, 18.43 mg EPA and 218.2 mg DHA, which is significantly more than in the 1st group, and also more than reported by the mentioned authors.

Modified diets administered to laying hens influenced the deposition of fatty acids in eggs. The EPA content was increased from 0.06% to 0.20%; the DHA content from 1.35% to 2.37% ($P < 0.001$). Compared to the 1st group, total n-3 PUFA were increased from 2.60% to 7.32% in total fatty acids in egg yolks of the 2nd group. Eggs laid in the 2nd group of hens contained 435 ALA, 18.4 EPA and 218.2 DHA mg/100 g of eggs, which marked an increase of x 4 for ALA, x 3 for

EPA, and x 1.7 for DHA in comparison to the 1st group. Furthermore, the total n-3 PUFA was increased in the 2nd group x 2.8 compared to the 1st group. This leads to the conclusion that consumption of eggs enriched with n-3 PUFA can have an important beneficial effect on human health.

Acknowledgement

This paper is a part of the research unit "Research, Production and Medical Testing of Functional Food" within "The Scientific Centre of Excellence for Personalized Health Care" financed by the Ministry of Science and Education of the Republic of Croatia.

References

- [1] Anonymous 1990. Nutrition Recommendations. Health and Welfare. Ministry of Supply and Services, Ottawa.
- [2] Antongiovanni MS, Minieri A, Buccioni I, Galligani I, Rapaccini S. 2009: Transfer of dietary fatty acid from butyric acid fortified canola oil into the meat of broiler. *Ital J Anim Sci* **8**: 754-756.
- [3] Aydin R, Dogan I 2010: Fatty acid profile and cholesterol content of egg yolk from chickens fed diets supplemented with purslane (*Portulaca oleraceae* L.). *J Sci Food Agric* **90**: 1759-1763.
- [4] Bavelaar FJ, Beynen AC 2004: Relationships between the intake of n-3 polyunsaturated fatty acids by hens and the fatty acid composition of their eggs. *International Journal of Poultry Science* **3**: 690-696.
- [5] Beynen AC 2004: Fatty acid composition of eggs produced by hens fed diets containing groundnut, soya bean or linseed. Fatty acid composition of eggs produced by hens fed diets containing groundnut, soya bean or linseed Fatty acid composition of eggs produced by hens fed diets containing groundnut, soya bean or linseed. *NJAS - Wagen J Life Sc* **52**: 3-10.
- [6] Csapó J, Sugár L, Horn A, Csapó Jne 1986: Chemical composition of milk from red deer, roe and fallow deer kept in captivity. *Acta Agronomica Hungarica* **36**: 359-372.
- [7] Ferrier LK, Caston LJ, Leeseon S, Squires J, Weaver BJ, Holub BJ 1995: Alpha-linolenic acid and docosahexaenoic acid enriched eggs from hens fed flaxseed: influence on blood lipids and platelet phospholipid fatty acids in humans. *Am J Clin Nutr* **62**: 81-86.
- [8] Gül M, Yoruk MA, Aksu T, Kaya A, Kaynar O 2012: The effect of different levels of canola oil on performance, egg shell quality and fatty acid composition of laying hens. *International Journal of Poultry Sciences* **11**: 769-776.
- [9] Hiltunen R, Holm Y, 2000: Lipid. In: Hiltunen, R. & Holm, Y. (eds.). *Farmakognosia. Yliopistopaino, Helsinki*. P. 142-170.
- [10] He X, Yang X, Guo Y 2007: Effects of different dietary oil sources on immune function in cyclophosphamide immunosuppressed chickens. *Anim Feed Sci Technol* **139**: 186-200.
- [11] Herkel R, Gálik R, Arpášova H, Bíro D, Juráček M, Šimko M, Rolinec M 2016: Fatty acid profile and nutritional composition of table eggs after supplementation by pumpkin and flaxseed oils. *Acta Vet Brno* **85**: 277-283.
- [12] Husveth F, Rozsa L, Magyar L, Bali G, Papocsi P 2003: N-3 fatty acid enrichment of table eggs by adding a fish oil preparation (Nordos Fat^R) to the diet of laying hens. *Arch Geflugelkd* **67**: 198-203.
- [13] Huyghebaert G, Raes K, Maertens L, Arnouts S, Delezie E. 2007: The interactive impact of dietary PUFA on the deposition of DHA in the egg yolk. In: *Proceedings of the 16th European Symposium on Poultry Nutrition*, in August in Strasbourg, France, p. 681-684.

- [14] Jiang Z, Ahn DU, Sim J 1991: Effect of feeding flax and two types of sunflower seeds on fatty acid compositions of yolk lipid classes. *Poult Sci* **41**: 2467-2475.
- [15] Kralik G, Škrčić Z, Gajčević Z, Hanžek D 2007: Utjecaj različitih ulja u hrani za nesilice na kvalitetu jaja i sadržaj masnih kiselina u žumanjku jajeta (Effect of different oils supplemented to laying hens' diets on the quality of eggs and the fatty acids content in egg yolk. *Krmiva* **49**: 115-125.
- [16] Kralik G, Gajčević Z, Škrčić Z 2008: The effect of different oil supplementations on laying performance and fatty acid composition of egg yolk. *Ital J Anim Sci* **7**: 173-183.
- [17] Lewis NM, Seburg S, Flanagan NL 2000: Enriched eggs a source of fatty acids for humans. *Poult Sci* **79**: 971-974.
- [18] Mantzioris E 2000: Biochemical effects of a diet containing food enriched with n-3 fatty acids. *Am J Clin Nutr* **72**: 42-48.
- [19] Meluzzi A, Sirri F, Manfreda G, Tallarico N, Franchini A 2000: Effect of dietary vitamin E on the quality of table eggs enriched with n-3 long chain fatty acids. *Poult Sci* **79**: 539-545.
- [20] Nykter M, Kymäläinen HR 2006: Quality characteristics of edible linseed oil. *Agri Food Sci* **15**: 402-413
- [21] Okuyama H, Kobayashi T, Watanabe S, 1997: Dietary fatty acids - the n-6/n-3 balance and chronic elderly disease. Excess linolenic acid and relative n-3 deficiency syndrome seen in Japan. *Prog Lipid Res* **35**: 409-457.
- [22] Pita MCG, Piber Neto E, Carvalho PR, Mendonca Junior CX 2010: Efeito da suplementação, óleo de canola e vitamina E na dieta sobre as concentrações de ácidos graxos poliinsaturados em ovos de galinha. *Arquivo Brasileiro De Medicina Veterinária E Zootecnia* **59**: 789-796 (in Portuguese).
- [23] Raes K, Huyghebaert G, De Smet S, Nollet L, Arnouts S, Demeyer D 2002: The deposition on conjugated linolenic acids in eggs of laying hens fed diets varying in fat level and fatty acid profile. *J Nutr* **132**: 182-189.
- [24] Rašić L, Čavka A, Kralik Z, Kralik G, Šerić V, Drenjančević I. 2014: Effect of omega-3 enriched food on blood pressure microcirculation and lipid profile in young healthy men. In: Abstract Book of the Third International Symposium on Hypertension ISHOP3, in November in Osijek, Croatia, p. 13.
- [25] Rowghani E, Arab M, Nazifi S, Bakhtiari Z 2007: Effect of canola oil on cholesterol and fatty acid composition of egg-yolk of laying hens. *International Journal of Poultry Science* **6**: 111-114.
- [26] Salamatdoustnobar R, Nazeradi K, Ayazi A, Hamidiyan A, Gorbani A, Fani, A 2009: Beneficial effects of canola oil on serum biochemical parameters of Iranian native turkeys. *J Anim Vet Adv* **8**: 2206-2209.
- [27] Samman S, Kung FP, Carter LM, Foster MJ, Ahmad ZI, Phuyal JL, Petocz P 2009: Fatty acid composition of certified organic, conventional and omega-3 eggs. *Food Chem* **116**: 911-914.
- [28] Sari M, Aksit M, Özdoğan M, Bamacioglu H, 2002: Effects of addition of flaxseed to diets of laying hens on some production characteristics, levels of yolk and serum cholesterol, and fatty acid composition of yolk. *Arch Geflügelkd* **66**: 75-79.
- [29] Scaife RJ, Moyo J, Galbraith H, Michie W, Campbell V 1994: Effect of dietary supplemental fats and oils on the tissue fatty acid composition and growth of female broilers. *Brit Poultry Sci* **35**: 107-118.
- [30] Scheideler SE 1997: Studies of consumer acceptance of high omega-3 fatty acid-enriched eggs. *J Appl Poultry Res* **6**: 137-146.
- [31] Simopoulos AP 2000: Human requirement for n-3 polyunsaturated fatty acids. *Poultry Sci* **79**: 961-970.

- [32] StatSoft, Inc. 2012. STATISTICA (data analysis software system), version 8.0. www.statsoft.com.
- [33] Sugano M 1996: Characteristics of fats in Japanese diets and current recommendations. *Lipids* **31**: 283-286.
- [34] Škrtić Z, Kralik G, Gajčević Z, Bogut I., Hanžek D 2007: The increase of the n-3 PUFA content in eggs. *Poljoprivreda/Agriculture/* **13**: 47-52.
- [35] Škrtić Z, Kralik G, Gajčević Z, Hanžek D 2008: Effect of different source of oils on fatty acid profile and organoleptic traits of eggs. *Acta agriculturae Slovenica* **92**: 213-217.
- [36] Van Elswik ME 1997: Comparison of n-3 fatty acids sources in laying hen rations for improvement of whole egg nutritional quality. *Brit J Nutr* **78**: S61-S69.
- [37] Yang CX, Ji C, Ding LM, Rong Y 2000: N-3 fatty acid metabolism and effects of alpha-linolenic acid on enriching n-3 FA eggs. *J Chi Agri Uni* **95**: 117-122.